



S35B2 Polyclonal Antibody

Catalog No	YP-Ab-06690
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	SLC35B2 PAPST1 PSEC0149
Protein Name	Adenosine 3'-phospho 5'-phosphosulfate transporter 1 (PAPS transporter 1) (Putative MAPK-activating protein PM15) (Putative NF-kappa-B-activating protein 48) (Solute carrier family 35 member B2)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	S35B2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	47kD
Cell Pathway	Golgi apparatus membrane ; Multi-pass membrane protein .
Tissue Specificity	Highly expressed in the placenta, pancreas, mammary gland and skeletal muscle. Weakly or not expressed in colon, heart and prostate.
Function	function:Mediates the transport of adenosine 3'-phospho 5'-phosphosulfate (PAPS), from cytosol into Golgi. PAPS is a universal sulfuryl donor for sulfation events that take place in the Golgi. PAPS indirectly participate in activation of the NF-kappa-B and MAPK pathways.,online information:GlycoGene database,similarity:Belongs to the nucleotide-sugar transporter family. SLC35B subfamily.,tissue specificity:Highly expressed in the placenta, pancreas, mammary gland and skeletal muscle. Weakly or not expressed in colon, heart and prostate.,
Background	Sulfotransferases (e.g., SULT4A1; MIM 608359) use an activated form of sulfate, 3-prime-phosphoadenosine 5-prime-phosphosulfate (PAPS), as a common sulfate donor for sulfation of glycoproteins, proteoglycans, and glycolipids in the endoplasmic reticulum and Golgi apparatus. SLC35B2 is located in the microsomal membrane and transports PAPS from the cytosol, where it is synthesized, into the Golgi lumen (Kamiyama et al., 2003 [PubMed



12716889)].[supplied by OMIM, Mar 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night