

C Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

SOX4 Polyclonal Antibody

and heart., Background SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member of the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Th encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (syntenin). The protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHrP) in bone development. The solution		
Reactivity Human;Mouse Applications WB;ELISA Gene Name SOX4 Protein Name Transcription factor SOX-4 Immunogen Synthesized peptide derived from part region of human protein Specificity SOX4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Nucleus . Tissue Specificity Testis, brain, and heart. Function Function function: Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAG-3' motif_similarity:Contains 1 MMG box DNA-binding domain_subunit:Interacts with UBE21, tissue specificity: Testis, brain and heart. Background SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member of the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic syndex of so transcription in grotein in, The enolded protein may	Catalog No	YP-Ab-06700
Applications WB;ELISA Gene Name SOX4 Protein Name Transcription factor SOX-4 Immunogen Synthesized peptide derived from part region of human protein Specificity SOX4 Polycional Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polycional, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Nucleus . Tissue Specificity Tissue Specificity Testis, brain, and heart. Function function: Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5:-AGAAAG-3 mattif, similarity: Contains 1 HMG box DNA-binding domainsubunit:Interacts with UBE21.,tissue specificity:Testis, brain and heart. Function SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member o the SOX (SRY-related HMG-box) family of transcription factor sinvolved in the endermination of the cell fate. Th encoded protein may duct as a transcriptional regulator dater forming a protein: complex with oth	Isotype	lgG
Gene NameSOX4Protein NameTranscription factor SOX-4ImmunogenSynthesized peptide derived from part region of human proteinSpecificitySOX4 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonyms-20°C/1 yearCell PathwayNucleus .Tissue SpecificityTestis, brain, and heart.Functionfunction. Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif, similarity Contains 1 HMG box DNA-binding domain, subunit:Interacts with UBE2L, tissue specificity. Testis, brain and heart.BackgroundSRY-box 4(SOX4) Homo sapiens end the determination of the cell fate. Th encoded protein may act as a transcriptional archivator that binds with high affinity to the T-cell enhancer with other proteins, such as syndecan binding protein (contens). The encoded protein may act as a transcriptional archivator that bindia protein forwing a protein complex with other proteins, such as syndecan binding protein (contens). The encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (contens). The encoded protein may act as a transcriptional arceulator the cell fate. Th encoded protein may act as a transcriptional ate	Reactivity	Human;Mouse
Protein Name Transcription factor SOX-4 Immunogen Synthesized peptide derived from part region of human protein Specificity SOX4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 52kD Cell Pathway Nucleus . Tissue Specificity Testis, brain, and heart. Function function:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5-AACAAAG-3' motif, similarity.Contains 1 HMG box DNA-binding domain, subunit.Interacts with UBE21, tissue specificity:Testis, brair and heart. Background SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member o the SOX (SRY-related HMG-box) family of transcription factors involved in the encoder protein may toct as a transcriptional regulator after forming a protein complex with other proteins, such as syndecaub inding protein (syntemin). The proteins with other proteins, such as syndecaub	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein Specificity SOX4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 52kD Cell Pathway Nucleus . Tissue Specificity Testis, brain, and heart. Function function:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5-AACAAAG-3' motif, similarity:Contains 1 HMG box DNA-binding domain, subunt:Interacts with UBE21, tissue specificity: Testis, brair and heart, Background SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member o the SOX (SRY-related HMG-box) family of transcription factors involved in the encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, subunt:Interacts with UBE21, tissue specificity: Testis, brair and heart,	Gene Name	SOX4
Specificity SOX4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit.lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 52kD Cell Pathway Nucleus . Tissue Specificity Testis, brain, and heart. Function function:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5-AACAAG-3' motif, similarity:Contains 1 HMG box DNA-binding domain, subunit:Interacts with UBE2.lissue specificity: Testis, brair and heart. Background SRY-box 4(SOX4) Homo sapiens This intronlexing a protein complex with other proteins, such as syndecan binding protein (synthenin). The protein may dat as a transcriptional regulator after forming a protein complex with other protein may dat as a transcriptional regulator after forming a protein complex with other protein may mediate downstream effects of paratryroid hormone (PTH) and PTH-related protein (PTH)Pi in bone development. The solution	Protein Name	Transcription factor SOX-4
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 52kD Cell Pathway Nucleus . Tissue Specificity Testis, brain, and heart. Function function:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif, similarity:Contains 1 HMG box DNA-binding domain.subunit:Interacts with UB:21.tissue specificity:Testis, brair and heart., Background SRY-box 4(SOX4) Homo sagiens This intronless gene encodes a member o the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Th erotein may act as a transcriptional activator after forming a protein (omplex with other protein may act as a transcriptional regulator of parathyroid hormone (PTH) and PTH-related protein (PTHP) in bone development. The solution of the may function in the apoptisis pathway leading to cell death as well as to turorigenesis and may mediate downstream effects of parathyroid hormone	Immunogen	Synthesized peptide derived from part region of human protein
SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsObserved Band52kDCell PathwayNucleus .Tissue SpecificityTestis, brain, and heart.Functionfunction:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif, similarity:Contains 1 HMG box DNA-binding domain, subunti:Interacts with UBE21, tissue specificity:Testis, brair and heart.BackgroundSRY-box 4(SOX4) Homo sapiens SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Th erooded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (syntenin). The protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provide	Specificity	SOX4 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 52kD Cell Pathway Nucleus . Tissue Specificity Testis, brain, and heart. Function function:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif, similarity:Contains 1 HMG box DNA-binding domain.,subunit:Interacts with UBE21.,tissue specificity:Testis, brair and heart. Background SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member on the SOX (SRY-related HMG-box) family of transcription factors involved in the ergulation of embryonic development and in the determining a protein complex with other protein may act as a transcriptional regulator after forming a protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHP) in bone development. The solution	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 52kD Cell Pathway Nucleus . Tissue Specificity Testis, brain, and heart. Function function:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif, similarity:Contains 1 HMG box DNA-binding domain.,subunit:Interacts with UBE21, tissue specificity:Testis, brair and heart. Background SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member o the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of the cell fate. Th encoded protein may act as a transcriptional regulator after forming a protein (syntenin). The protein may act as a transcriptional regulator after forming a protein (syntenin). The protein may and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. Eprovide	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 52kD Cell Pathway Nucleus . Tissue Specificity Testis, brain, and heart. Function function:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAG-3' motif, similarity:Contains 1 HMG box DNA-binding domain.,subunit:Interacts with UBE21, tissue specificity:Testis, brair and heart., Background SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member o the SOX (SRY-related HMG-box) family of transcription factors involved in the recoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTH/P) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provide	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 52kD Cell Pathway Nucleus . Tissue Specificity Testis, brain, and heart. Function function:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif, similarity:Contains 1 HMG box DNA-binding domain.,subunit:Interacts with UBE2I.,tissue specificity:Testis, brair and heart., Background SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member of the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Th encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (syntenin). The protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provide	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 52kD Cell Pathway Nucleus . Tissue Specificity Testis, brain, and heart. Function function:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif.,similarity:Contains 1 HMG box DNA-binding domain.,subunit:Interacts with UBE2I.,tissue specificity:Testis, brair and heart. Background SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member of the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Th encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (syntenin). The protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provide	Concentration	1 mg/ml
Synonyms Observed Band 52kD Cell Pathway Nucleus . Tissue Specificity Testis, brain, and heart. Function function:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif.,similarity:Contains 1 HMG box DNA-binding domain.,subunit:Interacts with UBE21.,tissue specificity:Testis, brair and heart., Background SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member of the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. The encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (syntenin). The protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHrP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provide	Purity	≥90%
Observed Band52kDCell PathwayNucleus .Tissue SpecificityTestis, brain, and heart.Functionfunction:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif.,similarity:Contains 1 HMG box DNA-binding domain.,subunit:Interacts with UBE21.,tissue specificity:Testis, brair and heart.,BackgroundSRY-box 4(SOX4) Homo sapiens the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Th encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (syntenin). The protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHrP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provide	Storage Stability	-20°C/1 year
Cell PathwayNucleus .Tissue SpecificityTestis, brain, and heart.Functionfunction:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif.,similarity:Contains 1 HMG box DNA-binding domain.,subunit:Interacts with UBE2I.,tissue specificity:Testis, brain and heart.,BackgroundSRY-box 4(SOX4) Homo sapiens regulation of embryonic development and in the determination of the cell fate. Th encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHrP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provide	Synonyms	
Tissue SpecificityTestis, brain, and heart.Functionfunction:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif., similarity:Contains 1 HMG box DNA-binding domain., subunit:Interacts with UBE2I., tissue specificity:Testis, brain and heart.,BackgroundSRY-box 4(SOX4) Homo sapiens the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Th encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHrP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provide	Observed Band	52kD
Functionfunction:Transcriptional activator that binds with high affinity to the T-cell enhancer motif 5'-AACAAAG-3' motif.,similarity:Contains 1 HMG box DNA-binding domain.,subunit:Interacts with UBE2I.,tissue specificity:Testis, brain and heart.,BackgroundSRY-box 4(SOX4) Homo sapiens the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Th encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (syntenin). The protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHrP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provide	Cell Pathway	Nucleus .
 enhancer motif 5'-AACAAAG-3' motif.,similarity:Contains 1 HMG box DNA-binding domain.,subunit:Interacts with UBE2I.,tissue specificity:Testis, brain and heart., Background SRY-box 4(SOX4) Homo sapiens This intronless gene encodes a member of the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Th encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (syntenin). The protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHrP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provide 	Tissue Specificity	Testis, brain, and heart.
the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. Th encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (syntenin). The protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHrP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provide	Function	enhancer motif 5'-AACAAAG-3' motif.,similarity:Contains 1 HMG box DNA-binding domain.,subunit:Interacts with UBE2I.,tissue specificity:Testis, brain,
	Background	the SOX (SRY-related HMG-box) family of transcription factors involved in the regulation of embryonic development and in the determination of the cell fate. The encoded protein may act as a transcriptional regulator after forming a protein complex with other proteins, such as syndecan binding protein (syntenin). The protein may function in the apoptosis pathway leading to cell death as well as to tumorigenesis and may mediate downstream effects of parathyroid hormone (PTH) and PTH-related protein (PTHrP) in bone development. The solution structure has been resolved for the HMG-box of a similar mouse protein. [provided]



UpingBio technology Co.,Ltd

Avoid repeated freezing and thawing!

C Tel: 400-999-8863 💌 Email:UpingBio@163.com

Website: www.upingBio.com

matters needing attention

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images