

VMAC Polyclonal Antibody

Catalog No	YP-Ab-06811
Isotype	lgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	VMAC
Protein Name	Vimentin-type intermediate filament-associated coiled-coil protein
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	VMAC Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	<u> </u>
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 18kD
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year 18kD Cytoplasm . Colocalizes with vimentin-type intermediate filaments
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year 18kD Cytoplasm . Colocalizes with vimentin-type intermediate filaments Cervix,Thymus,
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year 18kD Cytoplasm . Colocalizes with vimentin-type intermediate filaments Cervix,Thymus, subcellular location:Colocalizes with vimentin-type intermediate filaments.,
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year 18kD Cytoplasm . Colocalizes with vimentin-type intermediate filaments Cervix,Thymus, subcellular location:Colocalizes with vimentin-type intermediate filaments., subcellular location:Colocalizes with vimentin-type intermediate filaments.,










Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4°over night