





## PIGR Polyclonal Antibody

carcinoma-associated protein TB Immunogen  Synthesized peptide derived from Specificity  PIGR Polyclonal Antibody detects Formulation  Liquid in PBS containing 50% gly Source  Polyclonal, Rabbit, IgG  Purification  The antibody was affinity-purified affinity-chromatography using epi Dilution  WB 1:500-2000 ELISA 1:5000-20  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  84kD  Cell Pathway  [Polymeric immunoglobulin recep membrane protein .; [Secretory contains to the protein .; [Secretory contains the protein .]	endogenous levels of protein.
Reactivity Human;Rat;Mouse  Applications WB;ELISA  Gene Name PIGR  Protein Name Polymeric immunoglobulin recept carcinoma-associated protein TB Immunogen Synthesized peptide derived from Specificity PIGR Polyclonal Antibody detects  Formulation Liquid in PBS containing 50% gly  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified affinity-chromatography using epi  Dilution WB 1:500-2000 ELISA 1:5000-20  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 84kD  Cell Pathway [Polymeric immunoglobulin recept membrane protein .; [Secretory of Tissue Specificity Bile,Liver,Milk,Plasma,Saliva,Sm.	part region of human protein endogenous levels of protein.
Applications  Gene Name Protein Name Polymeric immunoglobulin recept carcinoma-associated protein TB Immunogen Synthesized peptide derived from Specificity PIGR Polyclonal Antibody detects Formulation Liquid in PBS containing 50% gly Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified affinity-chromatography using epi Dilution WB 1:500-2000 ELISA 1:5000-20 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year  Synonyms Observed Band 84kD Cell Pathway [Polymeric immunoglobulin recept membrane protein .; [Secretory of Tissue Specificity Bile, Liver, Milk, Plasma, Saliva, Smither Street Tissue Specificity Bile, Liver, Milk, Plasma, Saliva, Smither Street Tissue Specificity	part region of human protein endogenous levels of protein.
Gene Name       PIGR         Protein Name       Polymeric immunoglobulin recept carcinoma-associated protein TB         Immunogen       Synthesized peptide derived from         Specificity       PIGR Polyclonal Antibody detects         Formulation       Liquid in PBS containing 50% gly         Source       Polyclonal, Rabbit, lgG         Purification       The antibody was affinity-purified affinity-chromatography using epi         Dilution       WB 1:500-2000 ELISA 1:5000-20         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       84kD         Cell Pathway       [Polymeric immunoglobulin receptor contents in [Secretory cont	part region of human protein endogenous levels of protein.
Protein Name       Polymeric immunoglobulin recept carcinoma-associated protein TB         Immunogen       Synthesized peptide derived from Specificity         PIGR Polyclonal Antibody detects         Formulation       Liquid in PBS containing 50% gly         Source       Polyclonal, Rabbit, lgG         Purification       The antibody was affinity-purified affinity-chromatography using epi         Dilution       WB 1:500-2000 ELISA 1:5000-20         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       84kD         Cell Pathway       [Polymeric immunoglobulin recep membrane protein .; [Secretory or membrane protein .]         Tissue Specificity       Bile, Liver, Milk, Plasma, Saliva, Sn	part region of human protein endogenous levels of protein.
carcinoma-associated protein TB Immunogen  Synthesized peptide derived from Specificity  PIGR Polyclonal Antibody detects Formulation  Liquid in PBS containing 50% gly Source  Polyclonal, Rabbit, IgG  Purification  The antibody was affinity-purified affinity-chromatography using epi Dilution  WB 1:500-2000 ELISA 1:5000-20  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  84kD  Cell Pathway  [Polymeric immunoglobulin recep membrane protein .; [Secretory contains to the protein .; [Secretory contains the protein .]	part region of human protein endogenous levels of protein.
Specificity       PIGR Polyclonal Antibody detects         Formulation       Liquid in PBS containing 50% gly         Source       Polyclonal, Rabbit, IgG         Purification       The antibody was affinity-purified affinity-chromatography using epi         Dilution       WB 1:500-2000 ELISA 1:5000-20         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       84kD         Cell Pathway       [Polymeric immunoglobulin recep membrane protein .; [Secretory or membrane protein .; [Secretory or membrane protein .; [Secretory or membrane protein .; Secretory or m	endogenous levels of protein.
Formulation       Liquid in PBS containing 50% gly         Source       Polyclonal, Rabbit, lgG         Purification       The antibody was affinity-purified affinity-chromatography using epi         Dilution       WB 1:500-2000 ELISA 1:5000-20         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       84kD         Cell Pathway       [Polymeric immunoglobulin recepmembrane protein .; [Secretory commembrane protein .; [Secretor	·
Source       Polyclonal, Rabbit, IgG         Purification       The antibody was affinity-purified affinity-chromatography using epi         Dilution       WB 1:500-2000 ELISA 1:5000-20         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       84kD         Cell Pathway       [Polymeric immunoglobulin recepmembrane protein .; [Secretory contemplation of the protein of the purity of th	erol, and 0.02% sodium azide.
Purification       The antibody was affinity-purified affinity-chromatography using epi         Dilution       WB 1:500-2000 ELISA 1:5000-20         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       84kD         Cell Pathway       [Polymeric immunoglobulin recepmembrane protein .; [Secretory commembrane protein .]	
affinity-chromatography using epi  Dilution WB 1:500-2000 ELISA 1:5000-20  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 84kD  Cell Pathway [Polymeric immunoglobulin recepmembrane protein .; [Secretory content of the content of t	
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Synonyms         Observed Band       84kD         Cell Pathway       [Polymeric immunoglobulin recepmembrane protein .; [Secretory commembrane protein .]	•
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 84kD  Cell Pathway [Polymeric immunoglobulin recepmembrane protein .; [Secretory contents of the content	000
Storage Stability -20°C/1 year  Synonyms  Observed Band 84kD  Cell Pathway [Polymeric immunoglobulin recepmembrane protein .; [Secretory contents of the conte	
Synonyms  Observed Band  Cell Pathway  [Polymeric immunoglobulin recep membrane protein .; [Secretory contains to the contains of the contains to the contains	
Observed Band  Cell Pathway  [Polymeric immunoglobulin recepmembrane protein .; [Secretory contents of the con	
Cell Pathway  [Polymeric immunoglobulin recep membrane protein .; [Secretory continued to the continued to t	
Tissue Specificity  Membrane protein .; [Secretory continued to the contin	
· · · ·	or]: Cell membrane ; Single-pass type I mponent]: Secreted .
	all intestine,
epithelial cells. The complex is the the apical surface. During this proextracellular (known as the secre	neric IgA and IgM at the basolateral surface of en transported across the cell to be secreted at cess a cleavage occurs that separates the cory component) from the transmembrane
Background  polymeric immunoglobulin receptor member of the immunoglobulin sur polymeric immunoglobulin molecuthe complex is then transported a surface. A significant association nephropathy and several SNPs in	not necessary for Ig binding.,similarity:Contains ke) domains.,



## UpingBio technology Co.,Ltd

€ Tel: 400-999-8863 🛎 Email:UpingBio@163.com



matters	needing
attentio	

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images		