





THOP1 Polyclonal Antibody

Catalog No	YP-Ab-06921
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	THOP1
Protein Name	Thimet oligopeptidase (EC 3.4.24.15) (Endopeptidase 24.15) (MP78)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	THOP1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year 75kD
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year 75kD Cytoplasm.
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year 75kD Cytoplasm. Brain,Lung,Placenta,Uterus, catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues.,cofactor:Binds 1 zinc ion per subunit.,cofactor:Binds 1 zinc ion.,function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	75kD Cytoplasm. Brain,Lung,Placenta,Uterus, catalytic activity:Preferential cleavage of bonds with hydrophobic residues at P1, P2 and P3' and a small residue at P1' in substrates of 5 to 15 residues.,cofactor:Binds 1 zinc ion per subunit.,cofactor:Binds 1 zinc ion.,function:Involved in the metabolism of neuropeptides under 20 amino acid residues long. Involved in cytoplasmic peptide degradation. Able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments.,similarity:Belongs to the peptidase M3 family.,subunit:Monomer., The protein encoded by this gene is a kininase that uses zinc as a cofactor. The encoded oligopeptidase cleaves cytosolic peptides, making them unavailable for display on antigen-presenting cells. This protein also cleaves neuropeptides under 20 aa in length and can degrade beta-amyloid precursor protein to



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 🛎 Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images