







CLC14 Polyclonal Antibody

Catalog No YP-Ab-06946 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name CLEC14A C14orf27 EGFR5 UNQ236/PRO269 Protein Name C-type lectin domain family 14 member A (Epidermal growth factor receptor 5) (EGFR-5) Immunogen Synthesized peptide derived from part region of human protein Specificity CLC14 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD Cell Pathway Membrane; Single-pass type I membrane protein .
Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name CLEC14A C14orf27 EGFR5 UNQ236/PRO269 Protein Name C-type lectin domain family 14 member A (Epidermal growth factor receptor 5) (EGFR-5) Immunogen Synthesized peptide derived from part region of human protein Specificity CLC14 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD
Applications WB;ELISA Gene Name CLEC14A C14orf27 EGFR5 UNQ236/PRO269 Protein Name C-type lectin domain family 14 member A (Epidermal growth factor receptor 5) (EGFR-5) Immunogen Synthesized peptide derived from part region of human protein Specificity CLC14 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD
Gene Name CLEC14A C14orf27 EGFR5 UNQ236/PRO269 Protein Name C-type lectin domain family 14 member A (Epidermal growth factor receptor 5) (EGFR-5) Immunogen Synthesized peptide derived from part region of human protein Specificity CLC14 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD
Protein Name C-type lectin domain family 14 member A (Epidermal growth factor receptor 5) (EGFR-5) Immunogen Synthesized peptide derived from part region of human protein Specificity CLC14 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD
Immunogen Synthesized peptide derived from part region of human protein
Specificity CLC14 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 53kD
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 53kD
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms 53kD
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 53kD
Storage Stability -20°C/1 year Synonyms Observed Band 53kD
Synonyms Observed Band 53kD
Observed Band 53kD
Cell Pathway Membrane ; Single-pass type I membrane protein .
Tissue Specificity Blood,Brain,
Function similarity:Contains 1 C-type lectin domain.,similarity:Contains 1 EGF-like domain.,
This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. Members of this family share a common protein fold and have diverse functions, such as cell adhesion, cell-cell signalling, glycoprotein turnover, and roles in inflammation and immune response. This family member plays a role in cell-cell adhesion and angiogenesis. It functions in filopodia formation, cell migration and tube formation. Due to its presence at higher levels in tumor endothelium than in normal tissue endothelium, it is considered to be a candidate for tumor vascular targeting. [provided by RefSeq, Jan 2012],
matters needing Avoid repeated freezing and thawing! attention



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 🛎 Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images