



VWCE Polyclonal Antibody

Catalog No	YP-Ab-07006
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	VWCE URG11
Protein Name	von Willebrand factor C and EGF domain-containing protein (HBV X protein up-regulated gene 11 protein) (HBxAg up-regulated gene 11 protein)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	VWCE Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	105kD
Cell Pathway	Secreted . According to PubMed:16496348 is localized in the cytoplasm. .
Tissue Specificity	Expressed in liver.
Function	function:May be a regulatory element in the beta-catenin signaling pathway and a target for chemoprevention of hepatocellular carcinoma.,induction:By HBxAg.,similarity:Contains 4 EGF-like domains.,similarity:Contains 6 VWFC domains.,subcellular location:According to PubMed:16496348 is localized in the cytoplasm.,tissue specificity:Expressed in liver.,
Background	function:May be a regulatory element in the beta-catenin signaling pathway and a target for chemoprevention of hepatocellular carcinoma.,induction:By HBxAg.,similarity:Contains 4 EGF-like domains.,similarity:Contains 6 VWFC domains.,subcellular location:According to PubMed:16496348 is localized in the cytoplasm.,tissue specificity:Expressed in liver.,
matters needing attention	Avoid repeated freezing and thawing!



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images