

Ø Website: www.upingBio.com

H6ST2 Polyclonal Antibody

Isotype IgG Reactivity Human;Mouse Applications WB;ELISA Gene Name HS6ST2 PSEC0092 Protein Name Heparan-sulfate 6-O-sulfotransferase 2 (HS6ST-2) (EC 2.8.2) Immunogen Synthesized peptide derived from human protein . at AA range: 140-220 Specificity H6ST2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity< ≥90% 20°C/1 year Synonyms -20°C/1 year Synonyms -20°C/1 year Synonyms Catlalytic activity.3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3',5' bisphosphale + [heparan sulfate]-glucosamin		
Reactivity Human;Mouse Applications WB;ELISA Gene Name HS6ST2 PSEC0092 Protein Name Heparan-sulfate 6-O-sulfotransferase 2 (HS6ST-2) (EC 2.8.2) Immunogen Synthesized peptide derived from human protein . at AA range: 140-220 Specificity H6ST2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Gell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity.3°-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3°,5-bisphosphate + [heparan sulfate]-glucosamine = sulfate_intonion.6°,5-bisphosphate + [heparan sulfate]-glucosamine = adenosine 3°,5-bisphosphate + [heparan sulfate]-glucosamine = ad	Catalog No	YP-Ab-07242
Applications WB;ELISA Gene Name HS6ST2 PSEC0092 Protein Name Heparan-sulfate 6-O-sulfotransferase 2 (HS6ST-2) (EC 2.8.2) Immunogen Synthesized peptide derived from human protein . at AA range: 140-220 Specificity H6ST2 Polycional Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polycional, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity.3°-phosphoadenylyl sulfate + [heparan sulfate-glucosamine = adenosine 3'-5-bisphosphate + [heparan sulfate-glucosamine = denosine 3'-5-bisphosphate + [heparan sulfate-glucosamine = denosine 3'-5-bisphosphate + [heparan sulfate-glucosamine = adenosine 3'-5-bisphosphate + [heparan sulfate-glucosamine = denosine 3'-5-bisphosphate + [heparan sulfate-glucosamine = denosine 3'-5-bisphosphate + [heparan sulfate-glucosamine = denosine 3'	Isotype	lgG
Gene Name HS6ST2 PSEC0092 Protein Name Heparan-sulfate 6-O-sulfotransferase 2 (HS6ST-2) (EC 2.8.2) Immunogen Synthesized peptide derived from human protein . at AA range: 140-220 Specificity H6ST2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit.lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Call Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function Catalytic activity:3-phosphoadentyl sulfate + [heparan sulfate]-glucosamine = adenosine 3'.5-bisphosphate + [heparan sulfate]-glucosamine = defensine 3'.5-bisphosphate +	Reactivity	Human;Mouse
Protein Name Heparan-sulfate 6-O-sulfotransferase 2 (HS6ST-2) (EC 2.8.2) Immunogen Synthesized peptide derived from human protein . at AA range: 140-220 Specificity H6ST2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms G6kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3; 5'-bisphosphate + [heparan sulfate]-glucosamine = solloglucosamine residue (GlcNS) of heparan sulfate, similarity:Belongs to the sulfortansferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate (HS) Sulfotransferase gene family, which catalyze the transfer of sulfate	Applications	WB;ELISA
Immunogen Synthesized peptide derived from human protein . at AA range: 140-220 Specificity H6ST2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms G6kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = aderiosine 3; 5'-bisphosphate + [heparan sulfate]-glucosamine = sulfate, function:6-O-sulfation enzyme which catalyzes the transfer of sulfate from 3; biosphosphate + [heparan sulfate]-glucosamine = catalytic activity:3'-phosphoadenylyl sulfate + [Neparan sulfate]-glucosamine = aderiosine 3; 5'-bisphosphate (CloNS) of heparan sulfate, similarity:Belongs to the sulfortansferase 6 family. Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligande infinite to HS. Different family members and isofor	Gene Name	HS6ST2 PSEC0092
Specificity H6ST2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 66kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3°-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = actenosine 3°,5'-bisphosphate + [heparan sulfate]-glucosamine = sulfotoglucosamine residue (GICNS) of heparan sulfate, similarity:Belongs to the sulfotransferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate (HS) sulforansferase gene family, which catalyzes the transfer of sulfate to the solitorent isoforms have been and functions. Multiple transferior sulfate encoding different isoforms have been and functons. Multiple transferiory sulfate to HS). Dif	Protein Name	Heparan-sulfate 6-O-sulfotransferase 2 (HS6ST-2) (EC 2.8.2)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 66kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3,5'-bisphosphate + [heparan sulfate]-glucosamine = adenosine 3,5'-bisphosphate + [heparan sulfate]-glucosamine = sidenise from 3'-phosphoadenosine fibrion enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine fibrion enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine fibrion enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine fibrion enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine fibrion enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine fibrion enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine fibrion enzyme which catalyzes the transfer of sulfate fibrion enzyme which ca	Immunogen	Synthesized peptide derived from human protein . at AA range: 140-220
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 66kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3',5'-bisphosphate + [heparan sulfate]-glucosamine = sulfate, function: 6'-o-sulfation enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GIcNS) of heparan sulfate,similarity:Belongs to the sulformasferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, different family members and there the vith various ligands to influence cell growth, different family members and there the vith various ligands to influence cell growth, different family members and the fact with various ligands to influence cell growth, different family members and the soft with various ligands to influence cell growth, different family members and thex been	Specificity	H6ST2 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Observed Band 66kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3',5'-bisphosphate + (heparan sulfate]-glucosamine = 6-sulfate, function:6-O-sulfatio enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GlcNS) of heparan sulfate,similarity:Belongs to the sulfortransferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with usice-specific structure and functions. Multiple transfer of sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate to HS. Different family members and	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 66kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3',5'-bisphosphate + [heparan sulfate]-glucosamine = sulfotensine 3',5'-bisphosphoadensine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GIcNS) of heparan sulfate,similarity:Belongs to the sulforansferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfate swith tissue-specific structure and functions. Multiple transcript variants encoding different isoforms have been	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 66kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3',5'-bisphosphate + [heparan sulfate]-glucosamine = 6-sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GlcNS) of heparan sulfate.,similarity:Belongs to the sulfortansferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with tissue-specific structure and functions. Multiple transcript variants encoding different isoforms have been	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 66kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3',5'-bisphosphate + [heparan sulfate]-glucosamine = 6-sulfate.,function:6-O-sulfation enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GlcNS) of heparan sulfate,similarity:Belongs to the sulfotransferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate (HS) sulfotransferase gene family, which catalyze the transfer of sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with tissue-specific structure and functions. Multiple transcript variants encoding different isoforms have been	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band Observed Band 66kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3',5'-bisphosphate + [heparan sulfate]-glucosamine (6-sulfate, function:6-O-sulfation enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GlcNS) of heparan sulfate.,similarity:Belongs to the sulfotransferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate (HS) sulfotransferase gene family, which catalyze the transfer of sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with tissue-specific structure and functions. Multiple transcript variants encoding different isoforms have been	Concentration	1 mg/ml
Synonyms Observed Band 66kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3',5'-bisphosphate + [heparan sulfate]-glucosamine 6-sulfate.,function:6-O-sulfation enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GlcNS) of heparan sulfate.,similarity:Belongs to the sulfortansferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate (HS) sulfortansferase gene family, which catalyze the transfer of sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with tissue-specific structure and functions. Multiple transcript variants encoding different isoforms have been	Purity	≥90%
Observed Band 66kD Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3',5'-bisphosphate + [heparan sulfate]-glucosamine 6-sulfate.,function:6-O-sulfation enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GlcNS) of heparan sulfate.,similarity:Belongs to the sulfotransferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with tissue-specific structure and functions. Multiple transcript variants encoding different isoforms have been	Storage Stability	-20°C/1 year
Cell Pathway Membrane ; Single-pass type II membrane protein . Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine = adenosine 3',5'-bisphosphate + [heparan sulfate]-glucosamine 6-sulfate.,function:6-O-sulfation enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GlcNS) of heparan sulfate.,similarity:Belongs to the sulfotransferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate (HS) sulfotransferase gene family, which catalyze the transfer of sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with tissue-specific structure and functions. Multiple transcript variants encoding different isoforms have been	Synonyms	
Tissue Specificity Amygdala,Brain,Placenta, Function catalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine 6-sulfate.,function:6-O-sulfation enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GlcNS) of heparan sulfate.,similarity:Belongs to the sulfotransferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate (HS) sulfotransferase gene family, which catalyze the transfer of sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with tissue-specific structure and functions. Multiple transcript variants encoding different isoforms have been	Observed Band	66kD
Functioncatalytic activity:3'-phosphoadenylyl sulfate + [heparan sulfate]-glucosamine adenosine 3',5'-bisphosphate + [heparan sulfate]-glucosamine 6-sulfate.,function:6-O-sulfation enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GlcNS) of heparan sulfate.,similarity:Belongs to the sulfotransferase 6 family.,BackgroundHeparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with tissue-specific structure and functions. Multiple transcript variants encoding different isoforms have been	Cell Pathway	Membrane ; Single-pass type II membrane protein .
 adenosine 3',5'-bisphosphate + [héparan sulfate]-glucosamine 6 6-sulfate.,function:6-O-sulfation enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GlcNS) of heparan sulfate.,similarity:Belongs to the sulfotransferase 6 family., Background Heparan sulfate proteoglycans are ubiquitous components of the cell surface, extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate (HS) sulfotransferase gene family, which catalyze the transfer of sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with tissue-specific structure and functions. Multiple transcript variants encoding different isoforms have been 	Tissue Specificity	Amygdala,Brain,Placenta,
extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate (HS) sulfotransferase gene family, which catalyze the transfer of sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with tissue-specific structure and functions. Multiple transcript variants encoding different isoforms have been	Function	adenosine 3',5'-bisphosphate + [heparan sulfate]-glucosamine 6-sulfate.,function:6-O-sulfation enzyme which catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate (PAPS) to position 6 of the N-sulfoglucosamine residue (GlcNS) of heparan sulfate.,similarity:Belongs to the
	Background	extracellular matrix, and basement membranes, and interact with various ligands to influence cell growth, differentiation, adhesion, and migration. This gene encodes a member of the heparan sulfate (HS) sulfotransferase gene family, which catalyze the transfer of sulfate to HS. Different family members and isoforms are thought to synthesize heparan sulfates with tissue-specific structures and functions. Multiple transcript variants encoding different isoforms have been



Avoid repeated freezing and thawing!

🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com



matters needing attention

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images