






GPR21 Polyclonal Antibody

Catalog No	YP-Ab-07376
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	GPR21
Protein Name	Probable G-protein coupled receptor 21
Immunogen	Synthesized peptide derived from human protein . at AA range: 170-250
Specificity	GPR21 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year 38kD
Synonyms	·
Synonyms Observed Band	38kD
Synonyms Observed Band Cell Pathway	38kD Cell membrane; Multi-pass membrane protein. Not detected in the brain regions thalamus, putamen, caudate, frontal cortex,
Synonyms Observed Band Cell Pathway Tissue Specificity	38kD Cell membrane; Multi-pass membrane protein. Not detected in the brain regions thalamus, putamen, caudate, frontal cortex, pons, hypothalamus, hippocampus. function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Not detected in the brain regions thalamus, putamen,
Synonyms Observed Band Cell Pathway Tissue Specificity Function	Cell membrane; Multi-pass membrane protein. Not detected in the brain regions thalamus, putamen, caudate, frontal cortex, pons, hypothalamus, hippocampus. function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Not detected in the brain regions thalamus, putamen, caudate, frontal cortex, pons, hypothalamus, hippocampus., This gene encodes a member of the G-protein-coupled receptor 1 family. G-protein coupled receptors are membrane proteins which activate signaling cascades as a response to extracellular stress. The encoded protein activates a Gq signal transduction pathway which mobilizes calcium. [provided by RefSeq,
Synonyms Observed Band Cell Pathway Tissue Specificity Function Background	Cell membrane; Multi-pass membrane protein. Not detected in the brain regions thalamus, putamen, caudate, frontal cortex, pons, hypothalamus, hippocampus. function:Orphan receptor.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Not detected in the brain regions thalamus, putamen, caudate, frontal cortex, pons, hypothalamus, hippocampus., This gene encodes a member of the G-protein-coupled receptor 1 family. G-protein coupled receptors are membrane proteins which activate signaling cascades as a response to extracellular stress. The encoded protein activates a Gq signal transduction pathway which mobilizes calcium. [provided by RefSeq, Nov 2012],



Products Images



Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4° over night