




## TSK rabbit pAb

<b>Catalog No</b>	YP-Ab-07930
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	TSKU E2IG4 LRRC54 TSK UNQ850/PRO1788
<b>Protein Name</b>	TSK
<b>Immunogen</b>	Synthesized peptide derived from human TSK AA range: 288-338
<b>Specificity</b>	This antibody detects endogenous levels of TSK at Human/Mouse/Rat
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.44% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Tsukushin (Tsukushi) (E2-induced gene 4 protein) (Leucine-rich repeat-containing protein 54)
<b>Observed Band</b>	38kD
<b>Cell Pathway</b>	Secreted .
<b>Tissue Specificity</b>	Brain,Colon,Placenta,Uterus,
<b>Function</b>	induction:By 17-beta-estradiol.,similarity:Contains 9 LRR (leucine-rich) repeats.,
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images



Western blot analysis of lysates from VEC cells,  
primary antibody was diluted at 1:1000, 4° over night