


## EFCB7 rabbit pAb

Catalog No	YP-Ab-08253
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	EFCAB7 KIAA1799
Protein Name	EFCB7
Immunogen	Synthesized peptide derived from human EFCB7 AA range: 386-436
Specificity	This antibody detects endogenous levels of EFCB7 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentiation	i ing/iii
Purity	≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year  Cell projection, cilium membrane ; Peripheral membrane protein ; Cytoplasmic side . The EvC complex localizes at the base of cilia in the EvC zone of primary
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year  Cell projection, cilium membrane ; Peripheral membrane protein ; Cytoplasmic side . The EvC complex localizes at the base of cilia in the EvC zone of primary
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  Cell projection, cilium membrane ; Peripheral membrane protein ; Cytoplasmic side . The EvC complex localizes at the base of cilia in the EvC zone of primary cilia in a EFCAB7-dependent manner
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  Cell projection, cilium membrane ; Peripheral membrane protein ; Cytoplasmic side . The EvC complex localizes at the base of cilia in the EvC zone of primary cilia in a EFCAB7-dependent manner
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90%  -20°C/1 year  Cell projection, cilium membrane; Peripheral membrane protein; Cytoplasmic side. The EvC complex localizes at the base of cilia in the EvC zone of primary cilia in a EFCAB7-dependent manner  similarity:Contains 3 EF-hand domains.,

## **Products Images**



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night