







CI163 rabbit pAb


| Catalog No | YP-Ab-08257 |
|---|---|
| Isotype | IgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB |
| Gene Name | C9orf163 |
| Protein Name | CI163 |
| Immunogen | Synthesized peptide derived from human CI163 AA range: 144-194 |
| Specificity | This antibody detects endogenous levels of CI163 at Human |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1: 500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| C. man.ma | |
| Synonyms | |
| Observed Band | |
| - | |
| Observed Band | |
| Observed Band Cell Pathway | miscellaneous:This protein is produced by a bicistronic transcript which also produces the C9orf163 protein from an non-overlapping reading frame.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., |
| Observed Band Cell Pathway Tissue Specificity | miscellaneous:This protein is produced by a bicistronic transcript which also produces the C9orf163 protein from an non-overlapping reading frame.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., |
| Observed Band Cell Pathway Tissue Specificity Function | miscellaneous:This protein is produced by a bicistronic transcript which also produces the C9orf163 protein from an non-overlapping reading frame.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., Avoid repeated freezing and thawing! |
| Observed Band Cell Pathway Tissue Specificity Function Background matters needing | frame.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR., |







Products Images



Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night