






## NPAP1 rabbit pAb

| Catalog No                | YP-Ab-08279  |
|---------------------------|--|
| Isotype                   | IgG  |
| Reactivity                | Human;Rat;Mouse;   |
| Applications              | WB   |
| Gene Name                 | NPAP1 C15orf2  |
| Protein Name              | NPAP1  |
| Immunogen                 | Synthesized peptide derived from human NPAP1 AA range: 960-1010  |
| Specificity               | This antibody detects endogenous levels of NPAP1 at Human  |
| Formulation               | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source                    | Polyclonal, Rabbit,IgG   |
| Purification              | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  |
| Dilution                  | WB 1: 500-2000   |
| Concentration             | 1 mg/ml  |
| Purity                    | ≥90%   |
| Storage Stability         | -20°C/1 year   |
| Synonyms                  |  |
| Observed Band             |  |
| Cell Pathway              | Nucleus, nucleoplasm . Nucleus inner membrane . Colocalizes with the NPC and nuclear lamins at the nuclear periphery.  |
| Tissue Specificity        | Testis-specific in adults. In fetal brain expressed only from the paternal allele.   |
| Function                  | function:May be involved in spermatogenesis.,tissue specificity:Testis-specific in adults.,  |
| Background                | This gene is located in the Prader-Willi syndrome region on chromosome 15. This gene is biallelically expressed in adult testis and brain but is paternally imprinted in fetal brain. Defects in this gene may be associated with Prader-Willi syndrome. [provided by RefSeq, Aug 2012], |
| matters needing attention | Avoid repeated freezing and thawing!   |
| Usage suggestions         | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.  |
|                           |  |



## **Products Images**



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4°over night