


UFC1 rabbit pAb

Catalog No	YP-Ab-08688
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	UFC1 CGI-126 HSPC155
Protein Name	UFC1
Immunogen	Synthesized peptide derived from human UFC1 AA range: 90-140
Specificity	This antibody detects endogenous levels of UFC1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
•	
Storage Stability	-20°C/1 year
	-20°C/1 year
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year cytoplasm,extracellular exosome,
Storage Stability Synonyms Observed Band	
Storage Stability Synonyms Observed Band Cell Pathway	
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	cytoplasm,extracellular exosome, function:E2-like enzyme which forms an intermediate with UFM1 via a thioester linkage.,similarity:Belongs to the ubiquitin-conjugating enzyme family. UFC1
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	cytoplasm,extracellular exosome, function:E2-like enzyme which forms an intermediate with UFM1 via a thioester linkage.,similarity:Belongs to the ubiquitin-conjugating enzyme family. UFC1 subfamily., UFC1 is an E2-like conjugating enzyme for ubiquitin-fold modifier-1 (UFM1; MIM 610553) (Komatsu et al., 2004 [PubMed 15071506]).[supplied by OMIM, Mar
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	cytoplasm,extracellular exosome, function:E2-like enzyme which forms an intermediate with UFM1 via a thioester linkage.,similarity:Belongs to the ubiquitin-conjugating enzyme family. UFC1 subfamily., UFC1 is an E2-like conjugating enzyme for ubiquitin-fold modifier-1 (UFM1; MIM 610553) (Komatsu et al., 2004 [PubMed 15071506]).[supplied by OMIM, Mar 2008],



Products Images



Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night