







AN13B rabbit pAb


Catalog No	YP-Ab-08860
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	ANKRD13B
Protein Name	AN13B
Immunogen	Synthesized peptide derived from human AN13B AA range: 544-594
Specificity	This antibody detects endogenous levels of AN13B at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Cell membrane . Late endosome . Early endosome . Interaction with EGFR may
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Cell membrane . Late endosome . Early endosome . Interaction with EGFR may
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cell membrane . Late endosome . Early endosome . Interaction with EGFR may enhance association with the cell membrane. similarity:Contains 2 ANK repeats.,similarity:Contains 3 UIM (ubiquitin-interacting)
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cell membrane . Late endosome . Early endosome . Interaction with EGFR may enhance association with the cell membrane. similarity:Contains 2 ANK repeats.,similarity:Contains 3 UIM (ubiquitin-interacting)
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Cell membrane . Late endosome . Early endosome . Interaction with EGFR may enhance association with the cell membrane. similarity:Contains 2 ANK repeats.,similarity:Contains 3 UIM (ubiquitin-interacting motif) repeats.,







Products Images



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night