

🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com



## PGRP2 rabbit pAb

Catalog No	YP-Ab-09292
lsotype	lgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	PGLYRP2 PGLYRPL PGRPL UNQ3103/PRO10102
Protein Name	N-acetylmuramoyl-L-alanine amidase (EC 3.5.1.28) (Peptidoglycan recognition protein 2) (Peptidoglycan recognition protein long) (PGRP-L)
Immunogen	Synthesized peptide derived from human PGRP2 AA range: 90-140
Specificity	This antibody detects endogenous levels of human PGRP2
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Secreted. Membrane.
Cell Pathway Tissue Specificity	Secreted. Membrane. Strongly expressed in liver and fetal liver, and secreted into serum. Expressed to a much lesser extent in transverse colon, lymph nodes, heart, thymus, pancreas, descending colon, stomach and testis. Isoform 2 is not detected in the liver or serum.
	Strongly expressed in liver and fetal liver, and secreted into serum. Expressed to a much lesser extent in transverse colon, lymph nodes, heart, thymus, pancreas, descending colon, stomach and testis. Isoform 2 is not detected in the liver or
Tissue Specificity	Strongly expressed in liver and fetal liver, and secreted into serum. Expressed to a much lesser extent in transverse colon, lymph nodes, heart, thymus, pancreas, descending colon, stomach and testis. Isoform 2 is not detected in the liver or
Tissue Specificity Function	Strongly expressed in liver and fetal liver, and secreted into serum. Expressed to a much lesser extent in transverse colon, lymph nodes, heart, thymus, pancreas, descending colon, stomach and testis. Isoform 2 is not detected in the liver or
Tissue Specificity Function Background matters needing	Strongly expressed in liver and fetal liver, and secreted into serum. Expressed to a much lesser extent in transverse colon, lymph nodes, heart, thymus, pancreas, descending colon, stomach and testis. Isoform 2 is not detected in the liver or serum.



🔇 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

## **Products Images**