



# rabbit pAbFNIP2

<b>Catalog No</b>	YP-Ab-09420
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	FNIP2 FNIPL KIAA1450 MAPO1
<b>Protein Name</b>	Folliculin-interacting protein 2 (FNIP1-like protein) (O6-methylguanine-induced apoptosis 1 protein)
<b>Immunogen</b>	Synthesized peptide derived from human FNIP2 AA range: 745-795
<b>Specificity</b>	This antibody detects endogenous levels of human FNIP2
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:1000-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Lysosome membrane . Cytoplasm . Colocalizes with FLCN in the cytoplasm. .
<b>Tissue Specificity</b>	Widely expressed with highest levels in muscle, nasal mucosa, salivary gland, uvula, fat, liver, heart, placenta and pancreas (PubMed:18403135, PubMed:18663353, PubMed:27353360). Moderately expressed in the lung, small intestine, kidney and brain. Lower levels detected in renal cell carcinoma than in normal kidney tissue (PubMed:18403135). Higher levels detected in oncocytoma tumors than in normal kidney. Higher levels detected in renal cell carcinoma tumors than in normal kidney tissue (PubMed:27353360).
<b>Function</b>	
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**