



## NINL rabbit pAb

<b>Catalog No</b>	YP-Ab-09674
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	NINL KIAA0980 NLP
<b>Protein Name</b>	Ninein-like protein
<b>Immunogen</b>	Synthesized peptide derived from human NINL AA range: 1098-1148
<b>Specificity</b>	This antibody detects endogenous levels of human NINL
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:1000-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm . In interphase cells, NINL is transported to the centrosomes by the dynein-dynactin motor complex (PubMed:16254247). During centrosome maturation, PLK1 directly phosphorylates NINL resulting in its release into the cytoplasm (PubMed:16254247).
<b>Tissue Specificity</b>	Expressed in KYSE-150 esophageal carcinoma, HeLa cervical carcinoma and U2OS osteosarcoma cells. Expression is regulated in a cell cycle-dependent manner and peaks during G2/M phase (at protein level). Expressed in fetal heart, skeletal muscle, liver, lung and cochlea, and in adult brain, testis, kidney and retina.
<b>Function</b>	
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!



**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**