

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com





PI3R4 rabbit pAb

Catalog No	YP-Ab-09707	
Isotype	IgG	
Reactivity	Human;Mouse;Rat	
Applications	WB	
Gene Name	PIK3R4	
Protein Name	Phosphoinositide 3-kinase regulatory subunit 4 (PI3-kinase regulatory subunit 4) (EC 2.7.11.1) (PI3-kinase p150 subunit) (Phosphoinositide 3-kinase adaptor protein)	
Immunogen	Synthesized peptide derived from human PI3R4 AA range: 170-220	
Specificity	This antibody detects endogenous levels of human PI3R4	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.	
Source	Polyclonal, Rabbit,IgG	
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.	
Dilution	WB 1:1000-2000	
Concentration	4 mag/mal	
Concentration	1 mg/ml	
Purity	≥90%	
	<u> </u>	
Purity	≥90%	
Purity Storage Stability	≥90%	
Purity Storage Stability Synonyms	≥90%	
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Late endosome . Cytoplasmic vesicle, autophagosome . Membrane ; Lipid-anchor . As component of the PI3K complex I localized to pre-autophagosome structures. As component of the PI3K complex II localized predominantly to endosomes.	
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Late endosome . Cytoplasmic vesicle, autophagosome . Membrane ; Lipid-anchor . As component of the PI3K complex I localized to pre-autophagosome structures. As component of the PI3K complex II localized predominantly to endosomes. Localizes also to discrete punctae along the ciliary axoneme (By similarity)	
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Late endosome . Cytoplasmic vesicle, autophagosome . Membrane ; Lipid-anchor . As component of the PI3K complex I localized to pre-autophagosome structures. As component of the PI3K complex II localized predominantly to endosomes. Localizes also to discrete punctae along the ciliary axoneme (By similarity)	
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Late endosome . Cytoplasmic vesicle, autophagosome . Membrane ; Lipid-anchor . As component of the PI3K complex I localized to pre-autophagosome structures. As component of the PI3K complex II localized predominantly to endosomes. Localizes also to discrete punctae along the ciliary axoneme (By similarity)	
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Late endosome . Cytoplasmic vesicle, autophagosome . Membrane ; Lipid-anchor . As component of the PI3K complex I localized to pre-autophagosome structures. As component of the PI3K complex II localized predominantly to endosomes. Localizes also to discrete punctae along the ciliary axoneme (By similarity) Ubiquitously expressed.	



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



Products Images			