

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com





CECR1 rabbit pAb

	VP 41 00005	
Catalog No	YP-Ab-09805	
Isotype	IgG	
Reactivity	Human;Rat;Mouse;	
Applications	WB	
Gene Name	CECR1 ADA2 ADGF IDGFL	
Protein Name	Adenosine deaminase CECR1 (EC 3.5.4.4) (Cat eye syndrome critical region protein 1)	
Immunogen	Synthesized peptide derived from human CECR1 AA range: 243-293	
Specificity	This antibody detects endogenous levels of human CECR1	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.	
Source	Polyclonal, Rabbit,IgG	
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.	
Dilution	WB 1:1000-2000	
Concentration	1 mg/ml	
- "	>00%	
Purity	≥90%	
Storage Stability	-20°C/1 year	
Storage Stability		
Storage Stability Synonyms		
Storage Stability Synonyms Observed Band	-20°C/1 year	
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Secreted . Detected in blood plasma (at protein level). Widely expressed, with most abundant expression in human adult heart, lung, lymphoblasts, and placenta as well as fetal lung, liver, and kidney. In embryo, expressed in the outflow tract and atrium of the developing heart, the VII/VIII cranial nerve ganglion, and the	
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Secreted . Detected in blood plasma (at protein level). Widely expressed, with most abundant expression in human adult heart, lung, lymphoblasts, and placenta as well as fetal lung, liver, and kidney. In embryo, expressed in the outflow tract and atrium of the developing heart, the VII/VIII cranial nerve ganglion, and the	
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year Secreted . Detected in blood plasma (at protein level). Widely expressed, with most abundant expression in human adult heart, lung, lymphoblasts, and placenta as well as fetal lung, liver, and kidney. In embryo, expressed in the outflow tract and atrium of the developing heart, the VII/VIII cranial nerve ganglion, and the	
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	Secreted . Detected in blood plasma (at protein level). Widely expressed, with most abundant expression in human adult heart, lung, lymphoblasts, and placenta as well as fetal lung, liver, and kidney. In embryo, expressed in the outflow tract and atrium of the developing heart, the VII/VIII cranial nerve ganglion, and the notochord.	



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



Products Images			