



## VA0E2 rabbit pAb

<b>Catalog No</b>	YP-Ab-09863
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	ATP6V0E2 ATP6V0E2L C7orf32
<b>Protein Name</b>	V-type proton ATPase subunit e 2 (V-ATPase subunit e 2) (Lysosomal 9 kDa H(+)-transporting ATPase V0 subunit e2) (Vacuolar proton pump subunit e 2)
<b>Immunogen</b>	Synthesized peptide derived from human VA0E2 AA range: 26-75
<b>Specificity</b>	This antibody detects endogenous levels of human VA0E2
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:1000-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Membrane ; Multi-pass membrane protein . Cytoplasmic vesicle, clathrin-coated vesicle membrane ; Multi-pass membrane protein . Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane ; Multi-pass membrane protein .
<b>Tissue Specificity</b>	Isoform 1 is expressed at high levels in heart, brain and kidney and also detected in inner ear epithelium, vestibule, testis, epididymis and bladder. Isoform 2 is expressed in heart, kidney, placenta and pancreas. Isoform 2 is not detected in frontal cortex, but is prevalent in all other brain areas.
<b>Function</b>	
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images