

🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com

Website: www.upingBio.com

DOCK6 rabbit pAb

Catalog No	YP-Ab-09867
lsotype	lgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	DOCK6 KIAA1395
Protein Name	Dedicator of cytokinesis protein 6
Immunogen	Synthesized peptide derived from human DOCK6 AA range: 1460-1510
Specificity	This antibody detects endogenous levels of human DOCK6
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Cytoplasm . Cytoplasm, perinuclear region . Mainly located near the cell surface. Widely expressed. Expressed at low level in spleen, cerebellum, hippocampus
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cytoplasm . Cytoplasm, perinuclear region . Mainly located near the cell surface. Widely expressed. Expressed at low level in spleen, cerebellum, hippocampus
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cytoplasm . Cytoplasm, perinuclear region . Mainly located near the cell surface. Widely expressed. Expressed at low level in spleen, cerebellum, hippocampus
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Cytoplasm . Cytoplasm, perinuclear region . Mainly located near the cell surface. Widely expressed. Expressed at low level in spleen, cerebellum, hippocampus and in substantia nigra.



UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com



Products Images