

🔇 Tel: 400-999-8863 📼 Emall:Upingbio.163.com

Website: www.upingBio.com

## P4R3B rabbit pAb

Catalog No	YP-Ab-10008
Isotype	lgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	SMEK2 KIAA1387 PP4R3B PPP4R3B
Protein Name	Serine/threonine-protein phosphatase 4 regulatory subunit 3B (SMEK homolog 2
Immunogen	Synthesized peptide derived from human P4R3B AA range: 425-475
Specificity	This antibody detects endogenous levels of human P4R3B
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Observed Band Cell Pathway	Cytoplasm . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Nucleus . In interphase localized in the cytoplasm and (with higher levels) the nucleus. During metaphase located in pericentriolar regions.
	. Nucleus . In interphase localized in the cytoplasm and (with higher levels) the
Cell Pathway	. Nucleus . In interphase localized in the cytoplasm and (with higher levels) the nucleus. During metaphase located in pericentriolar regions.
Cell Pathway Tissue Specificity	. Nucleus . In interphase localized in the cytoplasm and (with higher levels) the nucleus. During metaphase located in pericentriolar regions.
Cell Pathway Tissue Specificity Function	. Nucleus . In interphase localized in the cytoplasm and (with higher levels) the nucleus. During metaphase located in pericentriolar regions.
Cell Pathway Tissue Specificity Function Background matters needing	. Nucleus . In interphase localized in the cytoplasm and (with higher levels) the nucleus. During metaphase located in pericentriolar regions. Moderately expressed in tissues and specific brain regions examined.



## UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com



**Products Images**