

🔇 Tel: 400-999-8863 📼 Emall:Upingbio.163.com

Website: www.upingBio.com

## TPC2A rabbit pAb

Catalog No	YP-Ab-10062
lsotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	TRAPPC2 SEDL
Protein Name	Trafficking protein particle complex subunit 2 (Sedlin)
Immunogen	Synthesized peptide derived from human TPC2A AA range: 45-95
Specificity	This antibody detects endogenous levels of human TPC2A
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
•	
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year Cytoplasm, perinuclear region . Endoplasmic reticulum-Golgi intermediate
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Cytoplasm, perinuclear region . Endoplasmic reticulum-Golgi intermediate compartment . Nucleus . Cytoplasm . Localized in perinuclear granular structures Expressed in brain, heart, kidney, liver, lung, pancreas, placenta, skeletal muscle
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Cytoplasm, perinuclear region . Endoplasmic reticulum-Golgi intermediate compartment . Nucleus . Cytoplasm . Localized in perinuclear granular structures Expressed in brain, heart, kidney, liver, lung, pancreas, placenta, skeletal muscle
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year Cytoplasm, perinuclear region . Endoplasmic reticulum-Golgi intermediate compartment . Nucleus . Cytoplasm . Localized in perinuclear granular structures Expressed in brain, heart, kidney, liver, lung, pancreas, placenta, skeletal muscle
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year Cytoplasm, perinuclear region . Endoplasmic reticulum-Golgi intermediate compartment . Nucleus . Cytoplasm . Localized in perinuclear granular structures Expressed in brain, heart, kidney, liver, lung, pancreas, placenta, skeletal muscle fetal cartilage, fibroblasts, placenta and lymphocytes.



## UpingBio technology Co.,Ltd

🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com



**Products Images**