

C Tel: 400-999-8863 ■ Emall:Upingbio.163.com





PASK (Phospho-Thr1165) rabbit pAb

Catalog No	YP-Ab-10513
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	PASK KIAA0135
Protein Name	PASK (Phospho-Thr1165)
Immunogen	Synthesized peptide derived from human PASK (Phospho-Thr1165)
Specificity	This antibody detects endogenous levels of PASK (Phospho-Thr1165) at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.169% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PAS domain-containing serine/threonine-protein kinase (PAS-kinase) (PASKIN) (hPASK) (EC 2.7.11.1)
Observed Band	
Cell Pathway	Cytoplasm . Nucleus . Localizes in the nucleus of testis germ cells and in the midpiece of sperm tails.
Tissue Specificity	Ubiquitously expressed, with slightly higher expression in brain, prostate and testis. Reduced expression was found in placenta. Present in germ cells of testis and in the midpiece of sperm tails (at protein level).
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Activated by autophosphorylation on Thr-1161 and Thr-1165 and inhibited by the first PAS domain.,PTM:Autophosphorylated on Thr-1161 and Thr-1165.,similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 PAS (PER-ARNT-SIM) domains.,tissue specificity:Ubiquitously expressed, with slightly higher expression in brain, prostate and testis. Reduced expression was found in placenta.,
Background	This gene encodes a member of the serine/threonine kinase family that contains two PAS domains. Expression of this gene is regulated by glucose, and the encoded protein plays a role in the regulation of insulin gene expression. Downregulation of this gene may play a role in type 2 diabetes. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this



Usage suggestions

UpingBio technology Co.,Ltd






gene. [provided by RefSeq, Nov 2011],

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of 1 Jurkat cell 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000