

(Tel: 400-999-8863 ■ Emall:Upingbio.163.com



Gz-α (Phospho-Ser16) rabbit pAb

| Catalog No | YP-Ab-10536 |
|---------------------------|---|
| Isotype | IgG |
| Reactivity | Human; Mouse;Rat |
| Applications | WB |
| Gene Name | GNAZ |
| Protein Name | Gz-α (Phospho-Ser16) |
| Immunogen | Synthesized peptide derived from human Gz-α (Phospho-Ser16) |
| Specificity | This antibody detects endogenous levels of $Gz-\alpha$ (Phospho-Ser16) at Human, Mouse,Rat |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.192% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | Guanine nucleotide-binding protein $G(z)$ subunit alpha $(G(x)$ alpha chain) $(Gz\text{-alpha})$ |
| Observed Band | 41kD |
| Cell Pathway | Membrane; Lipid-anchor. |
| Tissue Specificity | Brain,PNS, |
| Function | function:Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems.,similarity:Belongs to the G-alpha family. G(i/o/t/z) subfamily.,subunit:G proteins are composed of 3 units; alpha, beta and gamma. The alpha chain contains the guanine nucleotide binding site., |
| Background | G protein subunit alpha z(GNAZ) Homo sapiens The protein encoded by this gene is a member of a G protein subfamily that mediates signal transduction in pertussis toxin-insensitive systms. This encoded protein may play a role in maintaining the ionic balance of perilymphatic and endolymphatic cochlear fluids. [provided by RefSeq, Jul 2008], |
| matters needing attention | Avoid repeated freezing and thawing! |
| | |



UpingBio technology Co.,Ltd






Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of SY-SY5Y using primary antibody at 1:1000 dilution 4°C, overnight. Secondary antibody(catalog#:RS23920) was diluted at 1:10000 25 °C, 1.5hours