

🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com 🛛 🙆 V

Website: www.upingBio.com

## Kizuna (Phospho-Thr379) rabbit pAb

Catalog No	YP-Ab-10555
Isotype	lgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	PLK1S1 C20orf19 KIZ NCRNA00153 HT013
Protein Name	Kizuna (Phospho-Thr379)
Immunogen	Synthesized peptide derived from human Kizuna (Phospho-Thr379)
Specificity	This antibody detects endogenous levels of Kizuna (Phospho-Thr379) at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.211% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
	5
Purity	≥90%
Purity Storage Stability	-
•	≥90%
Storage Stability	≥90% -20°C/1 year
Storage Stability Synonyms	<ul> <li>≥90%</li> <li>-20°C/1 year</li> <li>Centrosomal protein kizuna (Polo-like kinase 1 substrate 1)</li> <li>Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, cilium basal body . Localizes to centrosomes throughout the cell cycle. After centrosome duplication, it usually remains associated only with the mother centrosome, containing the older mature centriole and particles surrounding it. During prophase, additional particles accumulate around both</li> </ul>
Storage Stability Synonyms Observed Band	<ul> <li>≥90%</li> <li>-20°C/1 year</li> <li>Centrosomal protein kizuna (Polo-like kinase 1 substrate 1)</li> <li>Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, cilium basal body . Localizes to centrosomes throughout the cell cycle. After centrosome duplication, it usually remains associated only with the mother centrosome, containing the older mature centriole and particles surrounding it. During prophase, additional particles accumulate around both separating centrosomes. Does not accumulate at the microtubule minus ends, but instead localizes to the centrosomes and centrosome-surrounding area in a</li> </ul>
Storage Stability Synonyms Observed Band Cell Pathway	<ul> <li>≥90%</li> <li>-20°C/1 year</li> <li>Centrosomal protein kizuna (Polo-like kinase 1 substrate 1)</li> <li>Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, cilium basal body . Localizes to centrosomes throughout the cell cycle. After centrosome duplication, it usually remains associated only with the mother centrosome, containing the older mature centriole and particles surrounding it. During prophase, additional particles accumulate around both separating centrosomes. Does not accumulate at the microtubule minus ends, but instead localizes to the centrosomes and centrosome-surrounding area in a microtubule-independent and dependent manner, respectively</li> </ul>
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	<ul> <li>≥90%</li> <li>-20°C/1 year</li> <li>Centrosomal protein kizuna (Polo-like kinase 1 substrate 1)</li> <li>Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, cilium basal body . Localizes to centrosomes throughout the cell cycle. After centrosome duplication, it usually remains associated only with the mother centrosome, containing the older mature centriole and particles surrounding it. During prophase, additional particles accumulate around both separating centrosomes. Does not accumulate at the microtubule minus ends, but instead localizes to the centrosomes and centrosome-surrounding area in a microtubule-independent and dependent manner, respectively</li> </ul>



## UpingBio technology Co.,Ltd


🔇 Tel: 400-999-8863 📼 Email:Upingbio.163.com

Website: www.upingBio.com

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of HeLa cell, 2, LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000