

**(** Tel: 400-999-8863 ■ Emall:Upingbio.163.com



## FAM48A Polyclonal Antibody

Catalog No	YP-Ab-10628
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	FAM48A C13orf19 FP757
Protein Name	family with sequence similarity 48, member A
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human FAM48A. AA range:231-280
Specificity	The antibody detects endogenous FAM48A protein
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000, ELISA 1:10000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	FAM48A C13orf19 FP757
Observed Band	85kD
Cell Pathway	Nucleus .
Tissue Specificity	Highly expressed in testis, moderately in brain and pituitary gland. Expressed in several fetal tissues, including lung, brain, thymus and kidney. Expression is down-regulated in malignant prostate tissues.
Function	function:Required for MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) activation during gastrulation. Required for down-regulation of E-cadherin during gastrulation by regulating E-cadherin protein level downstream from NCK-interacting kinase (NIK) and independently of the regulation of transcription by Fgf signaling and Snail.,similarity:Belongs to the FAM48 family.,subunit:Interacts with MAPK14.,tissue specificity:Highly expressed in testis, moderately in brain and pituitary gland. Expressed in several fetal tissues, including lung, brain, thymus and kidney. Expression is down-regulated in malignant prostate tissues.,
Background	function:Required for MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) activation during gastrulation. Required for down-regulation of E-cadherin during gastrulation by regulating E-cadherin protein level downstream from NCK-interacting kinase (NIK) and independently of the regulation of



## UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 🗷 Emall:Upingbio.163.com



transcription by Fgf signaling and Snail.,similarity:Belongs to the FAM48 family.,subunit:Interacts with MAPK14.,tissue specificity:Highly expressed in testis, moderately in brain and pituitary gland. Expressed in several fetal tissues, including lung, brain, thymus and kidney. Expression is down-regulated in malignant prostate tissues.,

matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of mouse-kidney cells using Antibody diluted at 1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000