




TIF-IA Polyclonal Antibody


Catalog No	YP-Ab-10800
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC
Gene Name	RRN3 TIFIA
Protein Name	RNA polymerase I-specific transcription initiation factor RRN3 (Transcription initiation factor IA) (TIF-IA)
Immunogen	Synthesized peptide derived from human TIF-IA Polyclonal
Specificity	This antibody detects endogenous levels of TIF-IA.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	RNA polymerase I-specific transcription initiation factor RRN3 (Transcription initiation factor IA) (TIF-IA)
Observed Band	70kD
Cell Pathway	Nucleus, nucleolus .
Tissue Specificity	Hippocampus,Kidney,Pooled,Spleen,
Function	function:Required for efficient transcription initiation by RNA polymerase I.,similarity:Belongs to the RRN3 family.,
Background	function:Required for efficient transcription initiation by RNA polymerase I.,similarity:Belongs to the RRN3 family.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western blot analysis of 3T3 lysate, antibody was diluted at 1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).