



TKNK rabbit pAb

Catalog No	YP-Ab-10963
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	TAC3 NKNB UNQ585/PRO1155
Protein Name	TKNK
Immunogen	Synthesized peptide derived from human TKNK AA range: 6-56
Specificity	This antibody detects endogenous levels of TKNK at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Secreted.
Tissue Specificity	
Function	developmental stage:In pregnancy, the expression of NKB is confined to the outer syncytiotrophoblast of the placenta, significant concentrations of NKB can be detected in plasma as early as week 9, and plasma concentrations of NKB are grossly elevated in pregnancy-induced hypertension and pre-eclampsia.,function:Tachykinins are active peptides which excite neurons, evoke behavioral responses, are potent vasodilators and secretagogues, and contract (directly or indirectly) many smooth muscles.,similarity:Belongs to the tachykinin family.,
Background	This gene encodes a member of the tachykinin family of secreted neuropeptides. The encoded preproprotein is proteolytically processed to generate the mature peptide, which is primarily expressed in the central and peripheral nervous systems and functions as a neurotransmitter. This peptide is the ligand for the neurokinin-3 receptor. This protein is also expressed in the outer syncytiotrophoblast of the placenta and may be associated with pregnancy-induced hypertension and pre-eclampsia. Mutations in this gene are



associated with normosmic hypogonadotropic hypogonadism. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Feb 2016],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HuvEc cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).