








PCOTH rabbit pAb

Catalog No	YP-Ab-11263
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF
Gene Name	PCOTH
Protein Name	PCOTH
Immunogen	Synthesized peptide derived from human PCOTH AA range: 14-64
Specificity	This antibody detects endogenous levels of PCOTH at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1: 50-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Observed Band Cell Pathway	Cytoplasm .
	Cytoplasm . Expressed in prostate and testis. Weakly or not expressed in other tissues. Overexpressed in prostate cancers.
Cell Pathway	Expressed in prostate and testis. Weakly or not expressed in other tissues. Overexpressed in prostate cancers. function:May be involved in growth and survival of prostate cancer cells through
Cell Pathway Tissue Specificity	Expressed in prostate and testis. Weakly or not expressed in other tissues. Overexpressed in prostate cancers. function:May be involved in growth and survival of prostate cancer cells through the TAF-lbeta pathway.,tissue specificity:Expressed in prostate and testis. Weakly
Cell Pathway Tissue Specificity Function	Expressed in prostate and testis. Weakly or not expressed in other tissues. Overexpressed in prostate cancers. function:May be involved in growth and survival of prostate cancer cells through the TAF-Ibeta pathway.,tissue specificity:Expressed in prostate and testis. Weakly
Cell Pathway Tissue Specificity Function Background matters needing	Expressed in prostate and testis. Weakly or not expressed in other tissues. Overexpressed in prostate cancers. function:May be involved in growth and survival of prostate cancer cells through the TAF-lbeta pathway.,tissue specificity:Expressed in prostate and testis. Weakly or not expressed in other tissues. Overexpressed in prostate cancers.,

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Tris-EDTA,pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200(4° overnight.3, Secondary antibody was diluted at 1:200(room temperature, 45min).