

🕻 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

CRYL1 rabbit pAb

Catalog No YP-Ab-11300 Isotype IgG Reactivity Human; Mouse;Rat Applications WB Gene Name CRYL1 CRY Protein Name CRYL1 Immunogen Synthesized peptide derived from human CRYL1 AA range: 50-100 Specificity This antibody detects endogenous levels of CRYL1 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons.similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family,tissue specificity/Widely expressed, with highest levels in liver and kidney. Background The uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catabyzes the dehydrogenase family.tissue specificity/Widely expressed, with hi		
Reactivity Human; Mouse;Rat Applications WB Gene Name CRYL1 CRY Protein Name CRYL1 Immunogen Synthesized peptide derived from human CRYL1 AA range: 50-100 Specificity This antibody detects endogenous levels of CRYL1 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit.lgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cytoplasm . Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons.,similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family, tissue specificity.Widely expressed, with highest levels in liver and kidney. Background The uronate cycle functions as an alternative glucose matabolic pathway. accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-quionate into dehydroquionate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme. ado i	Catalog No	YP-Ab-11300
Applications WB Gene Name CRYL1 CRY Protein Name CRYL1 Immunogen Synthesized peptide derived from human CRYL1 AA range: 50-100 Specificity This antibody detects endogenous levels of CRYL1 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit.lgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cytoplasm . Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons_similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family, tissue specificity.Widely expressed, with highest levels in liver and kidney. Function The uronate cycle functions as an alternative glucose metabolic pathway. accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-guionate into dehydro-l-guionate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme. add is inhibited by inorgenic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq., Jul 2008].	Isotype	lgG
Gene Name CRYL1 CRY Protein Name CRYL1 Immunogen Synthesized peptide derived from human CRYL1 AA range: 50-100 Specificity This antibody detects endogenous levels of CRYL1 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons.similarity:Belongs to the 3-hydroxyacy/-CoA dehydrogenase family, tissue specificity:Widely expressed, with highest levels in liver and kidney. Function sequence cole functions as an alternative glucose metabolic pathway, accounting for about 25% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-guionate into dentry-Louionate in the uronate cycle functions as an alternative glucose netabolic pathway, accounting for about 25% of daily glucose catabolism. The product of the gene catalyzes the dehydrogenation of L-guionate into dentry-Louionate in the lens of the eye. [provided by RefSeq, Jul 2008],	Reactivity	Human; Mouse;Rat
Protein Name CRYL1 Immunogen Synthesized peptide derived from human CRYL1 AA range: 50-100 Specificity This antibody detects endogenous levels of CRYL1 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cytoplasm . Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasonssimilarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family, tissue specificity.Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasonssimilarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family, tissue specificity.Widely expressed, with highest levels in liver and kidney. Background The uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-guionate into denydro-L-guionate in the uronate cycle function sen an alterna	Applications	WB
ImmunogenSynthesized peptide derived from human CRYL1 AA range: 50-100SpecificityThis antibody detects endogenous levels of CRYL1 at Human/Mouse/RatFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.DilutionWB 1: 500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsCytoplasm .Cell PathwayCytoplasm .Tissue SpecificityWidely expressed, with highest levels in liver and kidney.Functionsequence caution:Unknown reasons.similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.tissue specificity:Widely expressed, with highest levels in liver and kidney.BackgroundThe uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-guionate into dehydro-L-guionate in the iorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lenso of the eye. [provided by RefSed, Jul 2008], matters needing	Gene Name	CRYL1 CRY
Specificity This antibody detects endogenous levels of CRYL1 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons_similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.,tissue specificity:Widely expressed, with highest levels in liver and kidney. Background The uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate in the enzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008],	Protein Name	CRYL1
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasonssimilarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase familytissue specificity:Widely expressed, with highest levels in liver and kidney. Background The unonate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate in the dens of the eye. [provided by RefSeq, Jul 2008]. matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human CRYL1 AA range: 50-100
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Cytoplasm . Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons .similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase familytissue specificity:Widely expressed, with highest levels in liver and kidney. Background The uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catabolism. The prod	Specificity	This antibody detects endogenous levels of CRYL1 at Human/Mouse/Rat
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Cytoplasm . Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons.,similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.,tissue specificity.Widely expressed, with highest levels in liver and kidney. Background The uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate in the erabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate in the erabolic pathway, accounting for about 5% of daily glucose tabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate in the erabolic pathway, accounting for about 5% of daily glucose tabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate in the erabolic pathway, accounting for about 5% of daily glucose tabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate in the erabolic pathway, accounting for about 5% of daily glucose tabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate in the erabolic pathway, accou	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Cytoplasm . Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons .similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.tissue specificity:Widely expressed, with highest levels in liver and kidney. Background The uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate into dehydro_L-gulonate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band -20°C/1 year Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons.,similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.,tissue specificity:Widely expressed, with highest levels in liver and kidney., Background The uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate into dehydro-L-gulonate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band -20°C/1 year Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasonssimilarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.,tissue specificity:Widely expressed, with highest levels in liver and kidney. Background The uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate into dehydro-L-gulonate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Dilution	WB 1: 500-2000
Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band -20°C/1 year Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons.,similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.,tissue specificity:Widely expressed, with highest levels in liver and kidney. Background The uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-guionate into dehydro-L-guionate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons.,similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.,tissue specificity:Widely expressed, with highest levels in liver and kidney., Background The uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate into dehydro-L-gulonate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Cytoplasm . Tissue Specificity Widely expressed, with highest levels in liver and kidney. Function sequence caution:Unknown reasons.,similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.,tissue specificity:Widely expressed, with highest levels in liver and kidney., Background The uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate into dehydro-L-gulonate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell PathwayCytoplasm .Tissue SpecificityWidely expressed, with highest levels in liver and kidney.Functionsequence caution:Unknown reasons.,similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.,tissue specificity:Widely expressed, with highest levels in liver and kidney.,BackgroundThe uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate into dehydro-L-gulonate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008],matters needingAvoid repeated freezing and thawing!	Synonyms	
Tissue SpecificityWidely expressed, with highest levels in liver and kidney.Functionsequence caution:Unknown reasons.,similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.,tissue specificity:Widely expressed, with highest levels in liver and kidney.,BackgroundThe uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate into dehydro-L-gulonate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008],matters needingAvoid repeated freezing and thawing!	Observed Band	
Functionsequence caution:Unknown reasons.,similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.,tissue specificity:Widely expressed, with highest levels in liver and kidney.,BackgroundThe uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate into dehydro-L-gulonate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008],matters needingAvoid repeated freezing and thawing!	Cell Pathway	Cytoplasm .
dehydrogenase family.,tissue specificity:Widely expressed, with highest levels in liver and kidney.,BackgroundThe uronate cycle functions as an alternative glucose metabolic pathway, accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate into dehydro-L-gulonate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008],matters needingAvoid repeated freezing and thawing!	Tissue Specificity	Widely expressed, with highest levels in liver and kidney.
 accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate into dehydro-L-gulonate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural role in the lens of the eye. [provided by RefSeq, Jul 2008], matters needing Avoid repeated freezing and thawing! 	Function	sequence caution:Unknown reasons.,similarity:Belongs to the 3-hydroxyacyl-CoA dehydrogenase family.,tissue specificity:Widely expressed, with highest levels in liver and kidney.,
	Background	accounting for about 5% of daily glucose catabolism. The product of this gene catalyzes the dehydrogenation of L-gulonate into dehydro-L-gulonate in the uronate cycle. The enzyme requires NAD(H) as a coenzyme, and is inhibited by inorganic phosphate. A similar gene in the rabbit is thought to serve a structural
	matters needing attention	Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

