

🕑 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

MOSC1 rabbit pAb

Catalog No	YP-Ab-11302
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;ELISA;IHC
Gene Name	MARC1 MOSC1
Protein Name	MOSC1
Immunogen	Synthesized peptide derived from human MOSC1 AA range: 156-206
Specificity	This antibody detects endogenous levels of MOSC1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
	≥90% -20°C/1 year
Purity Storage Stability Synonyms	
Purity Storage Stability	-20°C/1 year Mitochondrion outer membrane ; Single-pass type II membrane protein . Membrane ; Lipid-anchor . Mitochondrial import is mediated by AA 1-40 and
Purity Storage Stability Synonyms Observed Band	-20°C/1 year Mitochondrion outer membrane ; Single-pass type II membrane protein .
Purity Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Mitochondrion outer membrane ; Single-pass type II membrane protein . Membrane ; Lipid-anchor . Mitochondrial import is mediated by AA 1-40 and
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Mitochondrion outer membrane ; Single-pass type II membrane protein . Membrane ; Lipid-anchor . Mitochondrial import is mediated by AA 1-40 and requires ATP. cofactor:Molybdenum.,function:Probable oxidoreductase.,similarity:Contains 1
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year Mitochondrion outer membrane ; Single-pass type II membrane protein . Membrane ; Lipid-anchor . Mitochondrial import is mediated by AA 1-40 and requires ATP. cofactor:Molybdenum.,function:Probable oxidoreductase.,similarity:Contains 1
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year Mitochondrion outer membrane ; Single-pass type II membrane protein . Membrane ; Lipid-anchor . Mitochondrial import is mediated by AA 1-40 and requires ATP. cofactor:Molybdenum.,function:Probable oxidoreductase.,similarity:Contains 1 MOSC domain.,




🕑 Tel: 400-999-8863 💌 Email:UpingBio@163.com


Ø Website: www.upingBio.com

: www.upingBio.com

Products Images



Western blot analysis of lysates from 3T3 cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).