






BWR1B rabbit pAb

Catalog No	YP-Ab-11515
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	SLC22A18AS BWR1B BWSCR1B ORCTL2S SLC22A1LS
Protein Name	BWR1B
Immunogen	Synthesized peptide derived from human BWR1B AA range: 44-94
Specificity	This antibody detects endogenous levels of BWR1B at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
O a management in m	1 mg/ml
Concentration	1 mg/ml
Purity	≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90%
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Most abundantly expressed in gastrointestinal tissues. Expressed at lower levels
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Most abundantly expressed in gastrointestinal tissues. Expressed at lower levels in kidney and placenta. Expressed in fetal brain, liver, placenta, kidney and lung. tissue specificity:Most abundantly expressed in gastrointestinal tissues. Expressed at lower levels in kidney and placenta. Expressed in fetal brain, liver,
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Most abundantly expressed in gastrointestinal tissues. Expressed at lower levels in kidney and placenta. Expressed in fetal brain, liver, placenta, kidney and lung. tissue specificity:Most abundantly expressed in gastrointestinal tissues. Expressed at lower levels in kidney and placenta. Expressed in fetal brain, liver,
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Most abundantly expressed in gastrointestinal tissues. Expressed at lower levels in kidney and placenta. Expressed in fetal brain, liver, placenta, kidney and lung. tissue specificity:Most abundantly expressed in gastrointestinal tissues. Expressed at lower levels in kidney and placenta. Expressed in fetal brain, liver, placenta, kidneys and lung.,

Products Images



Western blot analysis of lysates from CACO2 cells, primary antibody was diluted at 1:1000, 4° over night