



# VIGLN rabbit pAb


<b>Catalog No</b>	YP-Ab-11596
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	HDLBP HBP VGL
<b>Protein Name</b>	VIGLN
<b>Immunogen</b>	Synthesized peptide derived from human VIGLN AA range: 370-420
<b>Specificity</b>	This antibody detects endogenous levels of VIGLN at Human/Mouse/Rat
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1: 500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm . Nucleus .
<b>Tissue Specificity</b>	
<b>Function</b>	function:Appears to play a role in cell sterol metabolism. It may function to protect cells from over-accumulation of cholesterol.,similarity:Contains 10 KH domains.,similarity:Contains 11 KH domains.,similarity:Contains 14 KH domains.,similarity:Contains 3 KH domains.,similarity:Contains 6 KH domains.,similarity:Contains 7 KH domains.,
<b>Background</b>	The protein encoded by this gene binds high density lipoprotein (HDL) and may function to regulate excess cholesterol levels in cells. The encoded protein also binds RNA and can induce heterochromatin formation. [provided by RefSeq, Mar 2016],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!



### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of lysates from Jarkat cells,  
primary antibody was diluted at 1:1000, 4° over night