







## SPT22 rabbit pAb

Catalog No	YP-Ab-11613
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;IHC
Gene Name	SPATA22
Protein Name	SPT22
Immunogen	Synthesized peptide derived from human SPT22 AA range: 196-246
Specificity	This antibody detects endogenous levels of SPT22 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
O = == = = = = = = = = = = = = = = = =	A manufact
Concentration	1 mg/ml
Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year  Chromosome . Localizes on meiotic chromosome axes. Accumulates on resected
Purity Storage Stability Synonyms Observed Band	≥90%
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90%  -20°C/1 year  Chromosome . Localizes on meiotic chromosome axes. Accumulates on resected DNA. Localization is dependent on MEIOB
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  Chromosome . Localizes on meiotic chromosome axes. Accumulates on resected DNA. Localization is dependent on MEIOB  Highly expressed in adult testis.
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  Chromosome . Localizes on meiotic chromosome axes. Accumulates on resected DNA. Localization is dependent on MEIOB  Highly expressed in adult testis.
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90%  -20°C/1 year  Chromosome . Localizes on meiotic chromosome axes. Accumulates on resected DNA. Localization is dependent on MEIOB  Highly expressed in adult testis.  tissue specificity:Highly expressed in adult testis.,











## **Products Images**



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).