


## CC079 rabbit pAb

| Catalog No  | YP-Ab-11678   |
|---|---|
| Isotype   | IgG   |
| Reactivity  | Human;Rat;Mouse;  |
| Applications  | IHC;IF  |
| Gene Name   | C3orf79   |
| Protein Name  | CC079   |
| Immunogen   | Synthesized peptide derived from human CC079 AA range: 5-55   |
| Specificity   | This antibody detects endogenous levels of CC079 at Human   |
| Formulation   | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                   |
| Source  | Polyclonal, Rabbit,IgG  |
| Purification  | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution  | IHC-p 1: 50-200. IF 1:50-200  |
|   |   |
| Concentration   | 1 mg/ml   |
| Concentration Purity  | 1 mg/ml<br>≥90%   |
|   | -   |
| Purity  | ≥90%  |
| Purity Storage Stability  | ≥90%  |
| Purity Storage Stability Synonyms   | ≥90%  |
| Purity Storage Stability Synonyms Observed Band   | ≥90%  |
| Purity Storage Stability Synonyms Observed Band Cell Pathway  | ≥90%  |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity                                     | ≥90%  |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function                            | ≥90%  |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing | ≥90% -20°C/1 year   |









## **Products Images**

Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 30min).