


TRI44 rabbit pAb

Catalog No	YP-Ab-11694
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	TRIM44 DIPB
Protein Name	TRI44
Immunogen	Synthesized peptide derived from human TRI44 AA range: 42-92
Specificity	This antibody detects endogenous levels of TRI44 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year intracellular,
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year intracellular, Highly expressed in testis. function:May play a role in the process of differentiation and maturation of neuronal cells.,similarity:Contains 1 B box-type zinc finger., This gene encodes a member of the tripartite motif (TRIM) family. The TRIM motif
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year intracellular, Highly expressed in testis. function:May play a role in the process of differentiation and maturation of neuronal cells.,similarity:Contains 1 B box-type zinc finger., This gene encodes a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, namely a RING, a B-box type 1 and a B-box
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year intracellular, Highly expressed in testis. function:May play a role in the process of differentiation and maturation of neuronal cells.,similarity:Contains 1 B box-type zinc finger., This gene encodes a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, namely a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. [provided by RefSeq, Jul 2008],



Products Images



Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4° over night