








IGLL5 rabbit pAb

Catalog No	YP-Ab-11745
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	IGLL5
Protein Name	IGLL5
Immunogen	Synthesized peptide derived from human IGLL5 AA range: 50-100
Specificity	This antibody detects endogenous levels of IGLL5 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Descrite :	≥90%
Purity	-0070
Storage Stability	-20°C/1 year
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Secreted .
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Secreted .
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year Secreted . Contrary to IGLL1, not expressed in pre-B-cells. This gene encodes one of the immunoglobulin lambda-like polypeptides. It is located within the immunoglobulin lambda locus but it does not require somatic rearrangement for expression. The first exon of this gene is unrelated to immunoglobulin variable genes; the second and third exons are the immunoglobulin lambda joining 1 and the immunoglobulin lambda constant 1 gene segments. Alternative splicing results in multiple transcript variants.
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background	-20°C/1 year Secreted . Contrary to IGLL1, not expressed in pre-B-cells. This gene encodes one of the immunoglobulin lambda-like polypeptides. It is located within the immunoglobulin lambda locus but it does not require somatic rearrangement for expression. The first exon of this gene is unrelated to immunoglobulin variable genes; the second and third exons are the immunoglobulin lambda joining 1 and the immunoglobulin lambda constant 1 gene segments. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2010],

Products Images



Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4° over night