

NIT1 rabbit pAb

Catalog No	YP-Ab-11754
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	NIT1
Protein Name	NIT1
Immunogen	Synthesized peptide derived from human NIT1 AA range: 32-82
Specificity	This antibody detects endogenous levels of NIT1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year
Synonyms	-20°C/1 year [Isoform 2]: Mitochondrion .; [Isoform 1]: Cytoplasm .
Synonyms Observed Band	
Synonyms Observed Band Cell Pathway	[Isoform 2]: Mitochondrion .; [Isoform 1]: Cytoplasm .
Synonyms Observed Band Cell Pathway Tissue Specificity	[Isoform 2]: Mitochondrion .; [Isoform 1]: Cytoplasm . Detected in heart, brain, placenta, liver, skeletal muscle, kidney and pancreas. sequence caution:Chimeric cDNA. Its C-terminal exon is derived from the last exon of the DEDD gene.,similarity:Belongs to the UPF0012 family.,similarity:Contains 1 CN hydrolase domain.,tissue specificity:Detected in
Synonyms Observed Band Cell Pathway Tissue Specificity Function	[Isoform 2]: Mitochondrion .; [Isoform 1]: Cytoplasm . Detected in heart, brain, placenta, liver, skeletal muscle, kidney and pancreas. sequence caution:Chimeric cDNA. Its C-terminal exon is derived from the last exon of the DEDD gene.,similarity:Belongs to the UPF0012 family.,similarity:Contains 1 CN hydrolase domain.,tissue specificity:Detected in heart, brain, placenta, liver, skeletal muscle, kidney and pancreas., This gene encodes a member of the nitrilase protein family with homology to bacterial and plant nitrilases, enzymes that cleave nitriles and organic amides to the corresponding carboxylic acids plus ammonia. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq,



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com

Website: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from PC-12 cells, primary antibody was diluted at 1:1000, 4° over night