








## KLD10 rabbit pAb

Catalog No	YP-Ab-11808
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB;IHC
Gene Name	KLHDC10 KIAA0265
Protein Name	KLD10
Immunogen	Synthesized peptide derived from human KLD10 AA range: 9-59
Specificity	This antibody detects endogenous levels of KLD10 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
•	
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  Nucleus . Cytoplasm .  developmental stage:Expressed in fetal brain, liver, lung, kidney and
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year  Nucleus . Cytoplasm .  developmental stage:Expressed in fetal brain, liver, lung, kidney and
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year  Nucleus . Cytoplasm .  developmental stage:Expressed in fetal brain, liver, lung, kidney and placenta.,similarity:Contains 6 Kelch repeats.,











## **Products Images**



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human meningioma. 1, Antibody was diluted at 1:200(4 ° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).