








TM192 rabbit pAb


Catalog No	YP-Ab-11844
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB;IHC
Gene Name	TMEM192
Protein Name	TM192
Immunogen	Synthesized peptide derived from human TM192 AA range: 27-77
Specificity	This antibody detects endogenous levels of TM192 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year
Synonyms	-20°C/1 year Lysosome membrane ; Multi-pass membrane protein . Late endosome .
Synonyms Observed Band	
Synonyms Observed Band Cell Pathway	Lysosome membrane ; Multi-pass membrane protein . Late endosome .
Synonyms Observed Band Cell Pathway Tissue Specificity	Lysosome membrane ; Multi-pass membrane protein . Late endosome . Strongly expressed in kidney, liver, lung and pancreas.
Synonyms Observed Band Cell Pathway Tissue Specificity Function	Lysosome membrane ; Multi-pass membrane protein . Late endosome . Strongly expressed in kidney, liver, lung and pancreas.
Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	Lysosome membrane; Multi-pass membrane protein. Late endosome. Strongly expressed in kidney, liver, lung and pancreas. PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the TMEM192 family.,



Products Images



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).