

## CTGE9 rabbit pAb

**IgG** Isotype

Human; Rat; Mouse; Reactivity

**WB Applications** 

**Gene Name** CTAGE9

CTGE9 **Protein Name** 

**Immunogen** Synthesized peptide derived from human CTGE9 AA range: 294-344

Specificity This antibody detects endogenous levels of CTGE9 at Human

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Polyclonal, Rabbit, IgG Source

**Purification** The antibody was affinity-purified from rabbit serum by affinity-chromatography

using specific immunogen.

**Dilution** WB 1: 500-2000

Concentration 1 mg/ml

≥90% Purity

Storage Stability -20°C/1 year

Synonyms

**Observed Band** 

Membrane; Single-pass membrane protein. **Cell Pathway** 

Tissue Specificity

disease:Autoantigen found several cancer types, including benign meningioma and cutaneous T-cell lymphoma (CTCL). Isoform 5A and isoform 5B are **Function** 

recognized by patients' sera with tumor specimen of various

origins.,disease:Tumor-associated antigen found in several cutaneous T-cell lymphoma (CTCL). Also found in colorectal and breast carcinomas, head and neck squamous cell carcinomas and melanoma.,function:Tumor-associated antigen., sequence caution: Translation N-terminally extended., similarity: Belongs to the cTAGE family., tissue specificity: At the protein level, isoform 5A is expressed only in testis. Other isoforms are expressed in several other normal

tissues, including brain, muscle and cranial skin. Isoform 5A (at the protein level) and isoform 5B are expressed in cutaneous T-cell lymphoma (CTCL) cell lines, colorectal carcinomas, breast carcinomas and melanoma. Isoform 5B, but not isof

**Background** 



## UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ■ Email:UpingBio@163.com




matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from SH-SY5Y cells, primary antibody was diluted at 1:1000, 4° over night