





TUSC1 rabbit pAb

| Catalog No | YP-Ab-11921 |
|---------------------------|--|
| Isotype | IgG |
| Reactivity | Human; Mouse |
| Applications | WB |
| Gene Name | TUSC1 |
| Protein Name | TUSC1 |
| Immunogen | Synthesized peptide derived from human TUSC1 AA range: 120-170 |
| Specificity | This antibody detects endogenous levels of TUSC1 at Human/Mouse |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1: 500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | |
| Cell Pathway | |
| Tissue Specificity | Widely expressed at low level. Expressed at higher level in testis, weakly expressed in muscle, colon, lung and spleen. Not detected in 3 non small cell lung carcinoma (NSCLC) cell lines with homozygous deletion of the 9p region, while it is down-regulated in 3 other tumor cell lines. |
| Function | tissue specificity:Widely expressed at low level. Expressed at higher level in testis, weakly expressed in muscle, colon, lung and spleen. Not detected in 3 non small cell lung carcinoma (NSCLC) cell lines with homozygous deletion of the 9p region, while it is down-regulated in 3 other tumor cell lines., |
| Background | This gene is located within the region of chromosome 9p that harbors tumor suppressor genes critical in carcinogenesis. It is an intronless gene which is downregulated in non-small-cell lung cancer and small-cell lung cancer cell lines, suggesting that it may play a role in lung tumorigenesis. [provided by RefSeq, Jul 2008], |
| matters needing attention | Avoid repeated freezing and thawing! |



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com

Website: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4° over night