


## KDEL2 rabbit pAb


| Catalog No                            | YP-Ab-11971  |
|---------------------------------------|--|
| Isotype                               | IgG  |
| Reactivity                            | Human;Rat  |
| Applications                          | WB;ELISA;IHC   |
| Gene Name                             | KDELC2 UNQ1904/PRO4350   |
| Protein Name                          | KDEL2  |
| Immunogen                             | Synthesized peptide derived from human KDEL2 AA range: 414-464   |
| Specificity                           | This antibody detects endogenous levels of KDEL2 at Human/Rat  |
| Formulation                           | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source                                | Polyclonal, Rabbit,IgG   |
| Purification                          | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.            |
| Dilution                              | WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000   |
| Concentration                         | 1 mg/ml  |
| Purity                                | ≥90%   |
| Storage Stability                     | -20°C/1 year   |
| Synonyms                              |  |
| Observed Band                         |  |
| Cell Pathway                          | Endoplasmic reticulum lumen .  |
|                                       |  |
| Tissue Specificity                    |  |
| Tissue Specificity Function           | similarity:Belongs to the KDELC family.,similarity:Contains 1 filamin repeat.,                                       |
|                                       | similarity:Belongs to the KDELC family.,similarity:Contains 1 filamin repeat.,                                       |
| Function                              | similarity:Belongs to the KDELC family.,similarity:Contains 1 filamin repeat.,  Avoid repeated freezing and thawing! |
| Function  Background  matters needing |  |



## **Products Images**



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).