







## OR9G1 rabbit pAb

Catalog No	YP-Ab-12021
Isotype	IgG

Human;Rat;Mouse; Reactivity

**WB** Applications

**Gene Name** OR9G1 OR9G5

OR9G1 **Protein Name** 

**Immunogen** Synthesized peptide derived from human OR9G1 AA range: 190-240

Specificity This antibody detects endogenous levels of OR9G1 at Human

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Polyclonal, Rabbit, IgG Source

The antibody was affinity-purified from rabbit serum by affinity-chromatography **Purification** 

using specific immunogen.

**Dilution** WB 1: 500-2000

Concentration 1 mg/ml

≥90% Purity

-20°C/1 year Storage Stability

Synonyms

**Observed Band** 

Cell membrane; Multi-pass membrane protein. **Cell Pathway** 

Tissue Specificity

function:Odorant receptor .,miscellaneous:The sequence shown here is derived from an EMBL/GenBank/DDBJ third party annotation (TPA) **Function** 

entry., similarity: Belongs to the G-protein coupled receptor 1 family.,

olfactory receptor family 9 subfamily G member 1(OR9G1) Homo sapiens **Background** 

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR)

arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone

receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by

RefSeq, Jul 2008],



## UpingBio technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from HuvEc cells, primary antibody was diluted at 1:1000, 4°over night