

🕑 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

## SASH1 rabbit pAb

Catalog No	YP-Ab-12061
Isotype	lgG
Reactivity	Human; Mouse
Applications	WB;IHC
Gene Name	SASH1 KIAA0790 PEPE1
Protein Name	SASH1
Immunogen	Synthesized peptide derived from human SASH1 AA range: 373-423
Specificity	This antibody detects endogenous levels of SASH1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm .
Tissue Specificity	Expressed ubiquitously, with highest levels in lung, placenta, spleen and thymus. Down-regulated in the majority (74%) of breast tumors in comparison with corresponding normal breast epithelial tissues. Expressed in the epidermis, epidermal keratinocytes, dermal fibroblasts and melanocytes (PubMed:23333244, PubMed:26203640).
Function	function:May have a role in a signaling pathway. Could act as a tumor suppressor.,similarity:Contains 1 SH3 domain.,similarity:Contains 2 SAM (sterile alpha motif) domains.,tissue specificity:Expressed ubiquitously, with highest levels in lung, placenta, spleen and thymus. Down-regulated in the majority (74%) of breast tumors in comparison with corresponding normal breast epithelial tissues.,
Background	
matters needing attention	Avoid repeated freezing and thawing!



## UpingBio technology Co.,Ltd

C Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**

1 KDa 180 ---140 --100 --75 --60 --45 --35 --25 --15 --10 --10 --

primary antibody was diluted at 1:1000, 4°over night

Western blot analysis of lysates from A431 cells,

Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).