







## CC134 rabbit pAb

Catalog No	YP-Ab-12070
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	CCDC134
Protein Name	CC134
Immunogen	Synthesized peptide derived from human CC134 AA range: 99-149
Specificity	This antibody detects endogenous levels of CC134 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Nucleus . Cytoplasm . Secreted . Endoplasmic reticulum . Accumulates in the nucleus in response to UV irradiation (PubMed:22644376)
Tissue Specificity	Expressed in cervical gland, cervical squamous epithelium, endometrium, stomach, kidney distal convoluted tubule, spermatogenic cells in testis, mammary gland, liver and striated muscle (at protein level) (PubMed:18087676, PubMed:23070808). Also detected in placenta (PubMed:18087676). Highest expression in testis relative to other tissues (PubMed:18087676). Detected in T cells and dendritic cells; highly expressed in activated CD8(+) T cells, and also expressed at lower levels in CD4(+) T cells (PubMed:25125657).
Function	similarity:Belongs to the UPF0388 family.,
Background	
matters needing attention	Avoid repeated freezing and thawing!



## UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com

Website: www.upingBio.com

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4°over night