

🕑 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

REXO4 rabbit pAb

Catalog No	YP-Ab-12194
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;IHC
Gene Name	REXO4 PMC2 XPMC2H
Protein Name	REXO4
Immunogen	Synthesized peptide derived from human REXO4 AA range: 312-362
Specificity	This antibody detects endogenous levels of REXO4 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
	· · · · · · · · · · · · · · · · · · ·
Purity	≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Nucleus, nucleolus . similarity:Belongs to the REXO4 family.,similarity:Contains 1 exonuclease domain.,subunit:Can bind ESR1 and ESR2. This interaction is abrogated by
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Nucleus, nucleolus . similarity:Belongs to the REXO4 family.,similarity:Contains 1 exonuclease domain.,subunit:Can bind ESR1 and ESR2. This interaction is abrogated by
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Nucleus, nucleolus . similarity:Belongs to the REXO4 family.,similarity:Contains 1 exonuclease domain.,subunit:Can bind ESR1 and ESR2. This interaction is abrogated by estrogen and augmented by tamoxifen treatment.,




UpingBio technology Co.,Ltd


🕲 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

Products Images



Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).