

🕻 Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

MT1H rabbit pAb

Catalog No	YP-Ab-12235
Isotype	lgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF
Gene Name	MT1H
Protein Name	MT1H
Immunogen	Synthesized peptide derived from human MT1H AA range: 6-56
Specificity	This antibody detects endogenous levels of MT1H at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1: 50-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	nucleus,cytoplasm,perinuclear region of cytoplasm,
Tissue Specificity	
Function	domain:Class I metallothioneins contain 2 metal-binding domains: four divalent ions are chelated within cluster A of the alpha domain and are coordinated via cysteinyl thiolate bridges to 11 cysteine ligands. Cluster B, the corresponding region within the beta domain, can ligate three divalent ions to 9 cysteines.,function:Metallothioneins have a high content of cysteine residues that bind various heavy metals; these proteins are transcriptionally regulated by both heavy metals and glucocorticoids.,similarity:Belongs to the metallothionein superfamily. Type 1 family.,subunit:Monomer.,
Background	
matters needing attention	Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 💌 Email:UpingBio@163.com

Ø Website: www.upingBio.com

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).